



**UNIVERSITI PUTRA MALAYSIA**

**LARVAL NUTRITION OF SILVER BARB BARBODES GONIONOTUS  
WITH AN EMPHASIS ON PROTEIN-ENERGY REQUIREMENTS AND  
FEEDING STRATEGY**

**CARINA MIRANDA TAYAG.**

**FP 2004 31**

**LARVAL NUTRITION OF SILVER BARB *BARBODES GONIONOTUS*  
WITH AN EMPHASIS ON PROTEIN-ENERGY REQUIREMENTS  
AND FEEDING STRATEGY**

**By**

**CARINA MIRANDA TAYAG**

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,  
in Fulfilment of the Requirements for the Degree of  
Master of Science**

**May 2004**



## DEDICATION

To my beloved parents, Carlito Tayag and Delinia Tayag, for the love, encouragement, prayers and for molding me through the years to become the person I am right now.

Abstract of the thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

**LARVAL NUTRITION OF SILVER BARB *BARBODES GONIONOTUS*  
WITH AN EMPHASIS ON PROTEIN-ENERGY REQUIREMENTS  
AND FEEDING STRATEGY**

**By**

**CARINA MIRANDA TAYAG**

**May 2004**

**Chairman: Associate Professor Dr. Mohd. Salleh Kamarudin, Ph.D.**

**Faculty: Agriculture**

A series of experiments were conducted to determine the possibility of rearing silver barb larvae on an artificial diet. Prior to the evaluation of artificial diet in rearing silver barb larvae, a study was conducted to determine the changes in the amino acid profile of silver barb from freshly stripped eggs to newly hatched larvae. The result showed that the amino acid pattern changed according to the developmental stages of the fish larvae.

The optimum protein requirement of silver barb larvae was determined using five isocaloric diets with varying protein levels (40, 45, 50, 55 and 60% protein). The results showed that the larval growth was highest when they were fed a diet containing 55% protein followed by those fed with 50, 45, 60 and 40% protein. Although the highest survival was obtained among larvae fed on a 50% protein diet, it was not significantly different ( $P>0.05$ ) from the survival of those fed on a 55% protein diet.

In the following study, the optimal protein-energy requirement of silver barb larvae was determined. Six experimental diets containing two levels of protein (50% and 55% protein) and three energy levels (18.8, 19.7 and 20.5 kJ g<sup>-1</sup>) were evaluated. In this study, larvae fed a diet containing 50% protein with 19.7 kJ g<sup>-1</sup> energy obtained the best growth and highest survival. This level was found to be optimum for growth and maintenance of the larvae.

An experiment was also conducted to find the best feeding rate for silver barb larvae. The present study showed a significant relationship between feeding rate and growth of the larvae. Based on growth, feed efficiency and survival, an optimal feeding rate of 45% BW d<sup>-1</sup> was found for silver barb larvae.

Lastly, this study revealed that silver barb larvae are visual feeder. Feeding the larvae throughout the day (day and night) did not give any beneficial effect for the larvae and resulted in lower growth and survival. This study suggested that feeding the larvae four times during daytime was the best feeding frequency for silver barb larviculture.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PEMAKANAN LARVA LAMPAM JAWA *BARBODES GONIONOTUS* DENGAN  
PENEKANAN TERHADAP KEPERLUAN PROTEIN-TENAGA  
DAN STRATEGI PEMBERIAN MAKANAN**

Oleh

**CARINA MIRANDA TAYAG**

Mei 2004

**Pengerusi: Profesor Madya Mohd. Salleh Kamarudin, Ph.D.**

**Faculti: Pertanian**

Beberapa siri eksperimen telah dilakukan untuk mengkaji kemungkinan menternak ikan lampam jawa menggunakan makanan buatan. Sebelum penilaian makanan buatan dalam pengkulturan larva lampam Jawa dilakukan, satu kajian telah dijalankan untuk mengetahui perubahan yang berlaku terhadap profil asid amino yang terkandung dalam telur yang baru dikeluarkan induk sehingga larva menetas. Kajian menunjukkan asid amino berubah mengikut tahap perkembangan larva ikan tersebut.

Keperluan protein optimum larva lampam Jawa ditentukan menggunakan lima diet isokalori dengan aras protein yang berbeza (40, 45, 50, 55 dan 60% protein). Keputusan menunjukkan bahawa pertumbuhan larva tertinggi didapati apabila larva diberi diet yang diberi mengandungi 55% protein dan diikuti larval yang diberi yang mengandungi 50, 45, 60 dan 40% protein. Walaubagaimanapun, larva yang diberi diet 50% protein menunjukkan kemandirian yang tertinggi tetapi tidak ketara berbeza ( $P>0.05$ ) dari larva yang diberi diet mengandungi 55% protein.

Keperluan protein-tenaga optimum bagi larva lampam Jawa ditentukan dalam kajian berikutnya. Enam diet ujian yang mengandungi dua aras protein (50% dan 55% protein) serta tiga aras tenaga (18.82, 19.66 dan 20.5 kJ g<sup>-1</sup>) telah diuji. Dalam kajian ini, larva yang diberi diet yang mengandungi 50% protein dengan 19.66 kJ g<sup>-1</sup> menunjukkan pertumbuhan terbaik serta kemandirian yang tertinggi. Aras ini didapati optimum untuk pertumbuhan dan penyenggaraan larva.

Satu eksperimen juga telah dilakukan untuk menentukan kadar pemberian makanan yang terbaik untuk larva lampam Jawa. Kajian menunjukkan hubungan ketara antara kadar pemberian makanan dengan pertumbuhan larva. Berdasarkan pertumbuhan, kecekapan makanan serta kemandirian, kadar pemberian makanan optimum bagi larva lampam Jawa adalah 45% berat badan hari<sup>-1</sup>.

Kajian ini menunjukkan bahawa larva lampam Jawa mencari makanan menggunakan penglihatan. Pemberian makanan kepada larva sepanjang hari (pada waktu siang dan malam) tidak memberikan sebarang kesan lebih baik terhadap larva malahan menyebabkan pertumbuhan dan kemandirian yang rendah. Kajian ini mencadangkan pemberian makanan sebanyak empat kali sehari pada waktu siang adalah lelerapan pemberian yang terbaik untuk pengkulturan larva lampam Jawa.

## ACKNOWLEDGEMENTS

"The Lord is my strength and my shield; my heart trusted in Him, and I am helped; therefore my heart greatly rejoices, and with my song I will praise Him" Psalm 28:7.

The completion of this thesis would have not been possible without the generous and helping hand of many individuals. There is no exact word to say to express how grateful I am for having them around.

I was truly indebted to my supervisor, my boss, and my mentor, Assoc. Prof. Dr. Mohd. Salleh Kamarudin, for all the support and encouragement which, provided me with much enthusiasm and motivation to make this paper possible. I am so grateful and proud to be one of the students to be under the supervision of one of the pillar and rock in the field of fish nutrition. I was also thankful for always welcoming me to accommodate all my inquiries even if he was about to leave for home.

I wanted to take this opportunity also to thank and acknowledge the support of these kindhearted individuals, Dr. Che Roos Saad and Mr. Aizam Zainal Abidin, members of my supervisory committee for helping me out and supervising especially when my supervisor was in Australia. I will always be grateful for those valuable consultation hours<sup>and</sup> opening their door for me

Just like the car, this study would not run and would not reach its intended location unless it contains enough fuel to support its needs. In regards to this, my appreciation goes to the Government of Malaysia for the research fund under the Intensification of Research in Priority Areas (IRPA) grant no. 01-02-04-0501.

To Dr. Dzolkifli for giving the permission to use his laboratory for the proximate analysis of my feed ingredients. My warm appreciation for the accommodation you gave to me even though the laboratory was always busy and crowded with students doing analysis just like me.

To Mr. Halim and Ibrahim in the Animal Nutrition lab for assisting me in conducting the proximate analysis.

To Mr. Ismail of the Jabatan Sains Haiwan, Malaysia for helping me in the amino acid analysis of my sample. I truly acknowledge the effort he did in order for my analysis to complete though sometimes or most of the time I have caused him headache and I have taken some of his break times just to finish the analysis. Sometimes I go to his laboratory with not enough chemicals for the analysis but he always tells me not to worry and he will look for the chemicals I needed.

To Dr. Anwar Abdul Rahim, I express my profound gratitude for sharing his expertise in Statistics for the data analysis. Thanks are also due to Dr. Jalloh and Dr. Osman for the help they extended regarding the statistical analysis of the data.

To my colleagues and friends in the Aquatic Biotechnology Laboratory Tito Emil, Dr. Samad, Ladan, Hamid Khoda Bakshsh, Tan, Lee, Mohammad, and Pharvaneh, for the helping hand they extended in order for this study to be completed.



To Samad and Ladan for all the technical help and for all the advices which motivates me to strive more. To the former, thanks for the effort in helping me to induce my experimental fish to spawn, though sometimes the trial was not successful. To the latter, for all the sisterly advices I received. I truly acknowledge the friendship I have from this kindhearted couple.

To the staff of Aquatic Resources Technology Laboratory especially to Jasni and Wak Jamal for their valuable assistance. Words are not enough to express how grateful I am for having these people around. They were so kind enough to me and they were always willing to help especially in getting and choosing the broodstocks. They didn't even mind if they had to catch broodstocks during holidays so long as I can run the experiment and finish it on time.

To Mohammad, for always being there when we needed something on the laboratory.

To Lirong, one of my good friends here in UPM, for always opening the door of his car for me to get to the places I wanted to go. Thanks for driving me around UPM especially during the time when I needed to beat the deadline of GSO. Though most of the time I made him feels dizzy because I always forgot something to take with me so we need to go back and come back again to GSO just to settle the matters. I wouldn't forget the time when the bus service has been halted, he unselfishly extended his helping hand in sending me always to my supervisors office. He always tells me that he was just a text message away.

To Flora, one of the nice person I met here in Malaysia. Thanks for sharing me your sentiments, even your ups and downs. Thanks for those teary-eyed moments and for those wacky thing we had together which somehow makes us feels soothed and relieved from all those faulty experiences.

To Bernadette, my best friend for all the support and encouragement that I can be firm and strong enough to go and face all the trials and hardships in life. Thanks for always being there for me especially during the time that I was really down.

To all my friends and colleagues in CLSU, especially to Joan and Jen who were always there for me, even during our college life. I was truly blessed to have friends like them, who always comfort and tells me that everything happens for a reason that God takes away something or someone when He has something better to give. Thanks for all those encouraging words.

To my mentors in CLSU especially to Dr. Tereso Abella and Dr. Arsenia Cagauan, for encouraging me to pursue higher degree. I was truly indebted for all the knowledge and wisdom I learned which inspires me that someday I can be just like them.

To the Filipino Community in Serdang for all the parties, foods, jokes and stories we shared with each other. Having them around makes me feel at home, that somehow they can help to comfort me especially when boredom and homesickness attacks. I wanted to take this opportunity that even through this way I can show my appreciation to those people who were always at my side and we're always there for me when the sailing gets rough. To ate Hazel, for always being there since the very moment I stepped down my feet in this foreign land. I will be forever thankful for having her around which

make me feels that I have a sister of my own whom I can depend on. To Tita Ging, for all the friendly and motherly advices and words of encouragements. I wanted to thank her also for always bringing me around so I can get to know a lot of people, even those diplomats.

To William, thanks for making me always happy and spending his time talking to me. I really do appreciate his effort in trying to cheer me up. He will always be a friend to me and his name will forever be engraved in my heart.

To ate Sarah, for all the advices she had given on me. For always telling me to be happy all the time even during the moments when smiling seems to be the hardest thing to do. Thanks for all the prayers Ate.

To all the brethren at the Jesus is Lord Church, especially to Ate Lily and Rose for feeding me the right kind food I needed most when I felt hunger for the Word of God. Thanks for all the comfort and all the effort they did in order for me to gradually understand that God is working in my life. I am grateful also to Robert for all the prayers and encouragement which strengthened me that I can do anything as long as I remained faithful to God.

To my mommy Suping, for all the prayers and support she had extended on me financially and emotionally.

To my brothers, who became my inspiration to strive for more. I wanted to thank them for taking care of me and loving through the years and for always telling me that I will be forever their Princess.

Above all, I praise and worship the **Almighty God and our Lord Jesus Christ** for all the never-ending blessings which He showered down on me since the day I opened up my eyes into this world.

I certify that an Examination Committee met on 28<sup>th</sup> May 2004 to conduct the final examination of Carina Miranda Tayag on her Master of Science thesis entitled "Larval Nutrition of Silver Barb *Barbodes gonionotus* with an Emphasis on Protein-Energy Requirements and Feeding Strategy" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

**Abdul Razak Bin Alimon, Ph.D.**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Mustafa Kamal Abdul Satar, Ph.D.**


Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Annie Christianus, Ph.D.**

Lecturer  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Roshada binti Hashim, Ph.D.**

Professor  
Universiti Sains Malaysia  
(Independent Examiner)



---

**GULAM RUSUL RAHMAT ALI, Ph.D.**

Professor/Deputy Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date: 22 JUL 2004

This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirements for the degree of name of degree. The members of the Supervisory Committee are as follows:

**Assoc. Prof. Mohd. Salleh Kamarudin, Ph.D.**

Associate Professor  
Faculty of Agriculture  
Universiti Putra Malaysia  
(Chairman)

**Che Roos Saad, Ph.D.**

Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)

**Aizam Zainal Abidin**

Faculty of Agriculture  
Universiti Putra Malaysia  
(Member)




**AINI IDERIS, Ph.D.**

Professor/Dean  
School of Graduate Studies  
Universiti Putra Malaysia

Date : **16 AUG 2004**

## DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

  
CARINA MIRANDA TAYAG

Date: 31/7/04

## TABLE OF CONTENTS

	<b>Page</b>
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	v
ACKNOWLEDGEMENTS	vii
APPROVAL	xi
DECLARATION	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
LIST OF APPENDICES	xviii
LIST OF ABBREVIATIONS	xix
 <b>CHAPTER</b>	
 <b>I INTRODUCTION</b>	 <b>1</b>
Background of the study	1
Statement of the Problem	3
The Significance of the Study	6
Objectives	6
 <b>II LITERATURE REVIEW</b>	 <b>7</b>
Biology of <i>Barbodes gonionotus</i>	7
Nutrient Requirements of Fish Larvae	9
Proteins and Amino Acids	10
Dietary Lipids	12
Dietary Energy Requirements	13
Larval Fish Diet	15
Live Food for Larvae	18
Feeding Regime for fish larvae	20
 <b>III GENERAL METHODOLOGY</b>	 <b>24</b>
Location of the Study	24
Production of Larvae	24
Experimental Set-up	25
Sampling	25
Water Management	25
Proximate Analysis of Fish Larvae and Experimental Diet	26
Moisture Content	26
Ash Content	26
Total Protein	26
Total Lipids	26
Total Protein and Lipid in Experimental Diet	27
Amino acids	27

	Gross energy determination	27
	Diet Preparation	28
	Statistical Analysis	28
<b>IV</b>	<b>CHANGES IN THE AMINO ACID PROFILE OF EGGS AND NEWLY HATCHED LARVAE OF SILVER BARB <i>BARBODES GONIONOTUS</i> (BLEEKER)</b>	<b>29</b>
	Introduction	29
	Materials and Methods	30
	Amino Acid Analysis	31
	Statistical Analysis	31
	Results	32
	Description of Eggs and Larvae	32
	Amino Acid Content	32
	Discussion	35
<b>V</b>	<b>GROWTH, SURVIVAL AND BODY COMPOSITION OF SILVER BARB <i>BARBODES GONIONOTUS</i> (BLEEKER) LARVAE FED DIETS WITH VARYING PROTEIN LEVELS</b>	<b>38</b>
	Introduction	38
	Materials and Methods	39
	Experimental Diets	39
	Growth Trial	40
	Growth Parameters	42
	Chemical Analysis	42
	Statistical Analysis	42
	Results	43
	Discussion	46
<b>VI</b>	<b>DIETARY PROTEIN AND ENERGY REQUIREMENTS OF SILVER BARB <i>BARBODES GONIONOTUS</i> LARVAE</b>	<b>51</b>
	Introduction	51
	Materials and Methods	52
	Experimental Animals	52
	Experimental Diets	52
	Feeding Trial	54
	Proximate Analysis	54
	Statistical analysis	55
	Results	56
	Discussion	61
<b>VII</b>	<b>EFFECT OF DIFFERENT FEEDING LEVEL ON THE GROWTH AND SURVIVAL OF <i>BARBODES GONIONOTUS</i> LARVAE FED WITH FORMULATED DIETS</b>	<b>65</b>
	Introduction	65
	Materials and Methods	66
	Proximate analysis	67
	Statistical analysis	67
	Results	68
	Discussion	72

<b>VIII</b>	<b>EFFECT OF DIFFERENT FEEDING FREQUENCY ON THE GROWTH AND SURVIVAL OF <i>BARBODES GONIONOTUS</i> LARVAE</b>	<b>76</b>
	Introduction	76
	Materials and Methods	77
	Experiment 1	77
	Experiment 2	78
	Water quality management	78
	Growth and Survival	79
	Data analysis	79
	Results	80
	Growth and Survival of Fish Larvae	80
	Experiment 1	80
	Experiment 2	81
	Discussion	86
<b>IX</b>	<b>GENERAL DISCUSSIONS</b>	<b>90</b>
	Conclusions	95
	<b>BIBLIOGRAPHY</b>	<b>96</b>
	<b>APPENDICES</b>	<b>118</b>
	<b>BIODATA OF THE AUTHOR</b>	<b>129</b>
	Submitted Papers	130



## LIST OF TABLES

Table		Page
1	Amino acid (% dry weight) composition of freshly stripped eggs, fertilized eggs and newly hatched larvae of <i>Barbodes gonionotus</i> .....	33
2	A/E ratios* of the essential amino acid composition of eggs and newly hatched larvae of <i>Barbodes gonionotus</i> .....	34
3	Ingredients and proximate analysis of the experimental diets..	41
4	Survival (%), growth performance and feed efficiency of silver barb larvae fed the experimental diets with varying protein levels for 16 days .....	44
5	Proximate composition (% $\pm$ SEM) of silver barb larvae fed diets with varying dietary protein levels .....	46
6	Composition (% as fed basis) and proximate analysis of experimental diets with varying protein and energy levels .....	53
7	Survival, growth and feed efficiency of silver barb <i>B.gonionotus</i> larvae fed with varying protein and energy levels for a period of 16 days .....	58
8	Body compositions of silver barb <i>B. gonionotus</i> larvae fed diet with varying protein and energy levels for a period of 16 days	61
9	Composition of experimental diet for silver barb larvae .....	68
10	Survival, growth and feed efficiency of <i>B.gonionotus</i> larvae at different feeding levels.....	70
11	Body composition (% dry matter) of <i>B. gonionotus</i> larvae fed diet different feeding levels .....	72
12	Growth and feed efficiency of <i>B.gonionotus</i> larvae fed at different feeding frequencies for 16 days <sup>1,2</sup> (Expt. 1) .....	83
13	Growth and feed efficiency of <i>B. gonionotus</i> larvae fed throughout the day at different feeding frequencies after 16 days of rearing period (Expt. 2) .....	84

LIST OF FIGURES

Figure		Page
1	Second order polynomial regression analysis between weight (mg) and dietary protein for silver barb larvae .....	45
2	Relationship between dietary protein and energy on a) % length gain, b) % weight gain and c) % survival of <i>B. gonionotus</i> ....	59
3	Relationship between protein and energy on a) FCR and b) PER of silver barb <i>B. gonionotus</i> larvae.....	60
4	Growth of silver barb larvae fed at different feeding rates after 16 days of rearing period .....	71
5	Changes in mean body weight of <i>Barbodes gonionotus</i> larvae at 4-days interval fed at different feeding frequencies .....	85



LIST OF APPENDICES

Appendix		Page
1	Procedure for moisture analysis.....	118
2	Procedure for ash analysis.....	119
3	Total Protein Analysis .....	120
4	Method for lipid analysis.....	121
5	Procedure for protein analysis in feed.....	122
6	HCL Hydrolysis.....	123
7	Performic Acid Hydrolysis.....	124
8	Alkaline Hydrolysis.....	125
9	Amino acid composition of the protein sources in the diet....	126
10	Nutritional composition of diet ingredients.....	126
11	Water quality parameters measured during the conduct of the study (Chapter V) .....	127
12	Water quality parameters measured during the conduct of the study (Chapter VI).....	127
13	Water quality parameters measured during the conduct of the study (Chapter VII).....	127
14	Water quality parameters measured during the conduct of the study (Chapter VIII).....	128



## LIST OF ABBREVIATIONS

ANOVA	Analysis of Variance
AOAC	Association of Analytical Chemist
BSA	Bovine Serum Albumin
BW	Body weight
CPE	Carp pituitary extract
CRD	Complete randomized design
DMRT	Duncan's multiple range test
DO	Dissolve oxygen
Expt.	Experiment
FCR	Feed conversion ratio
g	Gram
kJ	Kilo joule
L	Liter
N	Normality
P:E	Protein-to-energy ratio
PER	Protein efficiency ratio
SAS	Statistical Analysis System
SGR	Specific growth rate
UPM	Universiti Putra Malaysia
YSI	Yellow Spring Instrument
µm	Micrometer
EAA	Essential Amino Acid
FAA	Free Amino Acid

## CHAPTER I

### INTRODUCTION

#### Background of the Study

In the last ten years, aquaculture has been the fastest growing primary industry worldwide, with 39.4 million tons produced in 1998 (Tacon and Forster, 2000). The fishery sector in Malaysia plays an important role in the national economy in terms of income, foreign exchange and employment (Abu Talib and Alias, 1997). It also supplies 60% of the total animal protein intake of the population. The demand of fish, as the main source of protein is expected to increase from an annual consumption of 630,000 tons to over 1,580,000 tons by the year 2010 (DOF, 1995). As the fishery products from the sea are not expected to increase any further, aquaculture will assume a greater role to increase fish supply.

*Barbodes gonionotus* is one of the important species of aquaculture in Bangladesh (Hussain et al., 2002), Thailand (Wee and Ngamsnae, 1987; Koedprang and Na-Nakorn, 2000), Indonesia (Pongthana et al., 1999) as well as in Malaysia (Soong, 1963). It was originally introduced from Indonesia for pond culture in Malaysia during 1950's (Soong, 1963). It contributes 17.5% of the total output from freshwater ponds with a production of 2,505 tons in 1992 (Kechik, 1995). However, a reduction on the volume produced was observed in year 2000 and it accounts for approximately 4% only of the total freshwater fish produced in Malaysia (FAO, 2002). This reduction might be due to the lack of information in culturing this fish as well as in larval rearing which was believed to be the major bottleneck in fish production. This fish was categorized under Cyprinidae, the biggest Family of freshwater fish and its local name is lampam

Jawa (Mohsin and Ambak, 1983). This fish can be cultured in seasonal (roadside ditches, backyard ponds and rice fields) and perennial (tanks and ponds) water bodies often in combination with other species of carps (Hussain et al., 2002). It grows to a marketable size within 8-12 months but matures at 4-6 months of age (Koedprang and Na-Nakorn, 2000).

Silver barb was found to be a potential pituitary donor in inducing other species of fish to spawn in captivity (Kottelat, 1998). With regard to this, the artificial propagation of this fish was extensively practiced and was established since 1967 (Tangtropiros et al., 1990). However, even though the induced spawning of this fish had been carried out for more than 3 decades, studies on the larval rearing of this fish was in idle until recently. Jahangard (2003) reared this larvae using live food and he further noted that this larvae can be reared on alternative live food such as nematodes. However, by merely feeding the larvae with live food is not enough to determine their nutritional needs. This enigma gave birth to the field of larval nutrition. In conjunction to this, a noble research had been generated to develop an artificial diet as an economic live food alternative. Moreover, Rosenlund et al. (1997) stated that microdiet offers the opportunity to introduced nutrients that are not present in the live food.

Long before this new field evolved, most researchers have focused in determining live foods suitable for fish larvae. Later on, they studied the possibility of rearing fish larvae on formulated diet to minimize or eliminate the use of live food. Since then, a lot of studies have been done and results are often contradicting from each other. Nutritional requirements and larval rearing either marine or freshwater fish had been carried out through the years. Their findings suggested that fish larvae cannot be reared on an artificial diet alone and that live food must be supplied at their early life

stages (Dabrowski and Glogowski, 1977; Kestemont and Stalmans, 1992; Lazo et al., 2000). However, these findings were in contrast to the results found by several researchers wherein they suggested that fish larvae could be reared on artificial diet alone without compromising their growth and survival (Dabrowski et al., 1985; Zambonino Infante and Cahu, 1994; Diaz et al., 1997; Cahu et al., 1998; Gisbert and Williot, 2002).

Silver barb is a cyprinid fish where common carp (*Cyprinus carpio*) belongs. Common carp larvae showed good performance when only fed artificial diet, indicating that live food can be completely eliminated on the larval rearing of this fish (Alami-Durente et al., 1991; Charlon and Bergot, 1994; Kestemont, 1995). With this finding, the feasibility of rearing silver barb larvae on an artificial diet alone and determining the feeding strategy has yet to be defined. This study would also be of help to mitigate the problems often encounter in hatchery such as high mortality and high production cost which mainly comes from the production of live food and later could expand the aquaculture industry of silver barb.

### **Statement of the Problem**

This millennium is faced with a challenge of producing more fish for the increasing population of the world, particularly in Southeast Asia. Aquaculture is widely recognized as viable and profitable enterprise worldwide. It will continue to grow and supply an increasingly larger percentage of fishery products consumed because the oceans, seas and rivers are inadequately managed by human and their yield is unpredictable (Lovell, 1988). With the increasing numbers of farmers shifting from agriculture into an aquaculture venture and with the modification of culture system, a

consistent supply of quality seedstock from the hatchery must be obtained to meet the demand of the farmers. However, the availability of quality seedstock has been one of the most critical factors in the commercial production of fin fishes.

Presently, the initial feeding of fish larvae in intensive condition depends on the supply of live prey such as rotifers and *Artemia*. In an attempt to increase the nutritional value of rotifers and *Artemia*, they are typically fed polyunsaturated fatty acid (PUFA) rich diets prior to being fed to larval fish. However, this practice further increased the production cost. The nauplii of *Artemia salina* are obtained from durable eggs collected in the natural environment and were sold at high price. Although *Artemia* certainly holds many benefits for larviculture, there are several disadvantages. First, *Artemia* is expensive. Second, if used as a sole food, it does not satisfy the nutritional requirements of larvae and consequently needs to be enriched at a considerable expense (Bernabe and Guissi, 1982). *Artemia* have resulted in high degree of abnormalities such as malpigmentation and incomplete eye migration in flatfish (Tonheim et al., 2000). The mortality rate of fish larvae in hatcheries is high; therefore fish hatcheries do not often achieve high production. Thus, special care has to be taken to ensure that the environment and nutritional conditions are optimal for the fish larvae.

Due to the increasing demand for food protein and the continuing acceptance of the consumers, there was an increased in the consumption of this fish. However, a sudden decreased on the production of silver barb was observed in the year 1992-2000 (Kechik, 1995; FAO, 2002). During those years, research on the larval rearing of this fish is limited and this maybe the major reason for the reduction of fish produced. This phenomenon also prompted researchers to study the techniques and methods in fish larval rearing.



The culture of *Barbodes gonionotus* had increased due to the increasing demand for protein food. One of the methods used to determine the specific nutritional requirements of the fish larvae is through the use of artificial diets comprising different levels of protein, lipids and other nutrient components.

With the increasing interest in the culture of this species, there is an urgent need for reliable, effective and efficient hatchery techniques to obtain a high and constant production and to provide the aquaculture industry with effective feeding strategies. With regards to this, research efforts must focus mainly on the formulation of appropriate feeding procedures, optimization and development of larval diet. The development of an artificial diet may constitute a cost effective alternative for the labor-intensive production of live food. The diet can be stored for a certain period of time and can be formulated to meet the specific nutritional requirements of the growing larvae. It is hypothesized that the development of an artificial diet for *Barbodes gonionotus* larvae may help in order to increase the production of this fish by reducing the mortality rate during the larval stages and to reduce the production cost.